

**The First Record of the Copepod *Caligus cordyla* Pillai, 1963 (Copepoda: Siphonostomatoida), Parasitic on Carangid Fish *Megalaspis cordyla* (Linnaeus, 1785) in Northwest of the Arab Gulf, Iraq**

**Khalidah S. Al-Niaeem<sup>1\*</sup>, Suzan A. Al-Azizz<sup>2</sup> & Fatimah H. Al-Ataby<sup>2</sup>**

<sup>1</sup>Department of Fisheries and Marine Resources, College of Agriculture, University of Basrah, Basrah, Iraq

<sup>2</sup>Department of Microbiology, College of Veterinary Medicine, University of Basrah, Basrah, Iraq

\*Corresponding author: kalidah\_salim@yahoo.com

**Abstract:** The copepod *Caligus cordyla* Pillai, 1963 was found attached on the gill filaments of one species of carangid fishes, *Megalaspis cordyla* (Linnaeus, 1758) which was collected from northwest of the Arab Gulf during the period from October 2013 till July 2014. This represents the first record of *C. cordyla* in the Iraqi territorial waters of the Arab Gulf. With this record, five specific *Caligus* species are so far known from fishes of Iraq, in addition to one unidentified *Caligus* species. *M. cordyla* is now considered as a new host for this parasite in the Arab Gulf. With this record, *M. cordyla* is considered as a host for three copepod species in Iraq as well as a host for four cestode species and one nematode species.

Keywords: *Caligus cordyla*, *Megalaspis cordyla*, Arab Gulf, Iraq.

## Introduction

Over 72.2 % of the earth surface is covered by water, and fishes are ubiquitous inhabitants of this ecosystem. There are 33,700 recognized fish species in the World (Froese & Pauly, 2018). The fishes are the most successful vertebrate group and play extremely important ecological role (Noga, 2010).

Diseases and parasites affecting fish hosts can cause high mortalities in both cultured and wild species fishes. Such losses are considered as one of the major barrier against threat to the expanding of fish industry (Overstreet, 1990). Parasitic crustaceans are common on fish hosts in coastal marine and brackish waters. There are three major groups of Crustacea which include fish parasites. These crustacean groups include Isopoda, Branchiura and Copepoda (Pillai, 1967).

Among the 332 marine fish species of Iraq, 56.8% of these species belong to the order Perciformes which includes 38 families of which the family Carrangidae is represented with 33 species (Ali et al., 2018). Despite this number of carangid species, the parasitic fauna of only six species were investigated (see Mhaisen et al., 2018). Al-Ataby (2012), Al-Ataby et al. (2012) and Al-Niaeem et al. (2013) were the only articles concerned with the copepodal fauna of only two (out of 33) carangid species in Iraq.

The present study was designed to investigate the parasitic copepods from the torpedo scad *Megalaspis cordyla* in the northwest part of the Arab Gulf within the Iraqi territorial marine waters.

## Materials and Methods

A total of 214 fish specimens of the carangid fish *M. cordyla* were examined for ectoparasites during the period from September 2010 to February 2011. They were captured by trawl net from the Iraqi marine waters, northwest Arab Gulf (latitudes 47° 30' to 48° 15', longitude 30° 50' to 30° 00'). The fishes were transported to the laboratory, and copepod parasites were removed from the gill filaments and kept in 70% ethanol.

Before dissection, the copepods were cleared in lactic acid by using the wooden slide method (Humes & Gooding, 1964). Measurements were made by using an ocular micrometer. Drawings were done by using a camera Lucida. Copepods were identified on the basis of their morphological features according to Pillai (1963, 1967) and Kabata (1979a, b). Two copepod specimens were sent to Prof. Dr. Geoffrey A. Boxshall, Department of Zoology, British Museum (Natural History), London on 8<sup>th</sup> August 2012 for confirmation of the identification.

## Results

### *Caligus cordyla* Pillai, 1963 (Figs. 1 & 2)

Number of copepods: Four females.

Number of measured specimens: Two females.

Number of infected fishes: 20.

Host: *Megalaspis cordyla* (Linnaeus, 1758) Family Carangidae.

Site of infection: Gills.

Locality: Northwest Arab Gulf within the Iraqi territorial waters.

Material deposition: Voucher specimens were deposited in the British Museum (Natural History), London under the accessions BMNH 2012-227-228.

Female: Body length 5.9 mm long with a carapace elegantly rounded, nearly equal in length and width, postero-median lobe hardly projecting beyond the lateral lobes which are curved to close the posterior sinuses. Genital segment enlarged with narrowed anterior area (10 × 8 mm).

Abdomen broad, broadly fused to genital complex. Antenna, 3-segmented, proximal segment small, second segment unarmed, distal segment with a long curved claw. Maxilliped, 3-segmented, proximal segment armed, middle segment unarmed, distal segment with basal seta.

Leg 1, long, vestigial endopod tipped with two setules, first segment of exopod with row of setules on posterior edge and small hyaline membrane. Leg 2, small with large plumose inner seta on posterior edge, both outer and medial edges of base with large marginal membrane. The present investigation indicates the first occurrence of *C. cordyla* in Iraq and the Arab Gulf.

## Discussion

About 2,000 species of parasitic arthropods have been described and the majority of which belong to the family Caligidae of the subclass Copepoda of the class Hexanauplia. The family Caligidae has come under intense scrutiny with the development of culturing fishes in sea cages, due to the most notorious pests affecting wild and cultured marine fish species (Lester & Hayward, 2006).

Among studies concerning caligid copepods in Iraq, Mhaisen et al. (2018) enlisted 13 species of the family Caligidae. These included three species of the genus *Annuretes* (Adday, 2013; Al-Hasson et al., 2014; Al-Hasson, 2015; Khamees & Adday, 2017), five identified *Caligus* species in addition to one unidentified *Caligus* species (Al-Daraji, 1995; Jori & Mohamad, 2008; Al-Ataby, 2012; Adday, 2013; Khamees & Adday, 2013; Venmathi Maran et al., 2014; Al-Hasson, 2015), three species of the genus *Hermilius* (Adday, 2013; Venmathi

Maran et al., 2014) and one species of the genus *Mappates* (Al-Hasson, 2015; Al-Niaeem et al., 2017).

According to a recent account on marine fish parasites of Iraq (Mhaisen et al., 2018), the carangid fish *M. cordyla* is so far infected with four cestode species (*Callitetrarhynchus gracilis*, *Callitetrarhynchus* sp., *Progrillotia* sp. and *Pseudogrillotia spratti*), one nematode species (*Philometra megalaspidis*) and three crustacean species (*Caligus cordyla*, *Lernanthropus corniger* and *L. indicus*).




Figure 1: *Caligus cordyla*, maxilliped and legs, scale bar= 0.05 mm.




Figure 2: *Caligus cordyla*, adult female, scale bar= 0.2 mm.

## References

- Adday, T.K. (2013). Parasitic crustaceans of some marine fishes of Basrah province, Iraq. Ph. D. Thesis, Coll. Agric., Univ. Basrah: 302 pp.
- Al-Ataby, F.H.A. (2012). Taxonomical and histopathological study of some carangid fishes parasites in Basrah province, Iraq. M. Sc. Thesis, Coll. Vet. Med., Univ. Basrah: 104 pp.
- Al-Ataby, F.H.; Al-Niaeem, K.S. & Al-Azizz, S.A. (2012). A new record of the parasitic copepod *Lernanthropus indicus* (Pillai, 1967) (Copepoda: Lernanthropidae) from carangid fishes in north-west Arabian Gulf, Iraq. *Egypt. J. Exp. Biol. (Zool.)*, 8 (2): 175-179.
- Al-Daraji, S.A.M. (1995). Taxonomical and ecological studies on the metazoan, parasites of some marine fishes of Khor Al-Zubair estuary, north-west of the Arabian Gulf. Ph. D. Thesis, Coll. Agric., Univ. Basrah: 182 pp.
- Al-Hasson, H.A.H. (2015). Taxonomical and pathological studies on parasites of some perciform fishes in Iraqi marine waters. M. S. Thesis, Coll. Vet. Med., Univ. Basrah: 162 pp.
- Al-Hasson, H.A.A.; Al-Azizz, S.A. & Al-Niaeem, K.S. (2014). First record of *Anuretes similis* Ho & Lin, 2000 (Siphonostomatoida: Caligidae) parasitic on sordid rubberlip *Plectorhinchus sordidus* (Perciformes: Haemulidae) from northwest of Arab Gulf, Iraq. *J. Int. Acad. Res. Multidiscipl.*, 2 (8): 326-333.
- Ali, A.H.; Adday, T.K. & Khamees, N.R. (2018). Catalogue of marine fishes of Iraq. *Biol. Appl. Environ. Res.*, 2 (2): 298-368.

- Al-Niaeem, K.S.; Al-Azizz, S.A. & Al-Ataby, F.H. (2013). The first record of the copepod *Lernanthropus corniger* Yamaguti, 1954 parasitizing two carangid fishes in northwest of the Arab Gulf, Iraq. *Ekologija*, 59 (2): 95-98.
- Al-Niaeem, K.S.; Al-Azizz, S.A. & Al-Hasson, H.A.H. (2017). The first record of *Mappates plataxus* Rangnekar, 1958 (Copepoda: Siphonostomatoida: Caligidae) parasitic on longfin batfish *Platax teira* (Pisces: Ehippidae) from marine waters of Iraq. *Biol. Appl. Environ. Res.*, 1 (2): 219-227.
- Froese, R. & Pauly, D. (eds.) (2018). FishBase. World Wide Web electronic publication. [www.fishbase.org](http://www.fishbase.org). (Version 06/ 2018).
- Humes, A.G. & Gooding, R.U. (1964). A method for studying the external anatomy of copepods. *Crustaceana*, 6 (3): 238-240.
- Jori, M.M. & Mohamad, E.T. (2008). The effect of *Hamatopeduncularia* sp. and *Caligus* sp. on some blood parameters of *Arius bilineatus* (Val., 1840). *Mar. Mesopot.*, 23 (2): 269-277.
- Kabata Z. (1979a). Parasitic Copepoda of British fishes. The Ray Society, London: 50-68.
- Kabata, Z. (1979b). Copepoda of Australian fishes, XII. Family Lernanthropidae. *Crustaceana*, 37 (2): 198-213.
- Khamees, N.R. & Adday, T.K. (2013). Occurrence of sea lice *Caligus epinepheli* Yamaguti, 1936 (Copepoda: Siphonostomatoida) on gills of *Nemipterus japonicus* (Bloch, 1775) from northwest of the Arabian Gulf. *Basrah J. Agric. Sci.*, 26 (1): 1-14.
- Khamees, N.R. & Adday, T.K. (2017). Record of two *Anuretes* species (Copepoda: Siphonostomatoida) from fishes of the Arab Gulf, off Iraq. *Basrah J. Agric. Sci.*, 30 (2): 16-26.
- Lester, R.J.G. & Hayward, C.J. (2006). Phylum Arthropoda. In: Woo, P.T.K. (ed.). *Fish diseases and disorders, Vol. 1: Protozoan and metazoan infections*, 2<sup>nd</sup> edition. CAB Int., Wallingford: 466-565.
- Mhaisen, F.T.; Ali, A.H. & Khamees, N.R. (2018). Marine fish parasitology of Iraq: A review and checklists. *Biol. Appl. Environ. Res.*, 2 (2): 231-297.
- Noga, E.J. (2010). *Fish disease: Diagnosis and treatment*, 2<sup>nd</sup> edition. Wiley-Blackwell, Ames, Iowa: 519 pp.
- Overstreet, R.M. (1990). Metazoan parasitic diseases: Introductory remarks. In: Perkins, F.O. & Cheng, T.C. (eds.). *Pathology in marine science*. Acad. Press, San Diego, California: 261-266.
- Pillai, N.K. (1963). Copepods parasitic on south Indian fishes: Family Anthosomidae. *J. Bombay Nat. Hist. Soc.*, 60: 655-670.
- Pillai, N.K. (1967). Copepods parasitic on Indian marine fishes: A review. *Proceedings of Symposium on Crustacea*. Cochin, 5: 1556-1680.
- Venmathi Maran, B.A.; Moon, S.Y.; Adday, T.K.; Khamees, N.R. & Myoung, J.G. (2014). Three new records of copepods (Siphonostomatoida) parasitic on marine fishes of Iraq, including the relegation of two species of *Lernanthropinus* to *Lernanthropinus temminckii* (von Nordmann, 1864). *Acta Parasitol.*, 59 (1): 139-152.